



## E-QUAL Learning Collaboratives

[Sepsis](#) | [Opioid](#) | [Stroke](#)

### 2020 Enrollment Open!

- [Opioid Wave III](#) - April 20
- [Stroke Wave I](#)<sup>New</sup> - May 5

Enroll now through the E-QUAL Portal (*left*).

Currently enrolled sites may also enroll through the E-QUAL Portal. You will be prompted within the portal. Look for a **green bar** at the top of your portal page and follow the instructions.

---

E-QUAL Initiative Portal

---

Questions?

Contact the [E-QUAL Team](#)  
Use the [Portal Access Guide](#)

Quality Tip of the Month





“Patients who present to the emergency department in opioid withdrawal are suffering and vulnerable to dangerous addiction harms if symptoms are not managed effectively. There are a multitude of non-opioid therapies that have been traditionally used to treat opioid withdrawal that can be used in combination to improve symptoms, but non-opioid treatments do not address the underlying problem, do not mitigate cravings, and often result in prolonged ED lengths of stay and return visits. Optimal ED management takes advantage of the occasion of opioid withdrawal to initiate the best therapy for opioid addiction: medication assisted treatment with buprenorphine.”

*Reuben Strayer, MD*

*Maimonides Medical Center*

## Frequently Asked Questions



[View the FAQ page](#) | [Download the Portal Access Guide](#)

If you have questions regarding your collaborative, please be sure to reference these resources. If your site's questions are not answered, please reach out to the [E-QUAL team](#).

Click [here](#) if you are having trouble viewing this message.

© 2020 American College of Emergency Physicians. All rights reserved.

ACEP Corporate Offices • 4950 W. Royal Lane • Irving, Texas 75063

Getting too many emails? Update your [ACEP Email Subscription Center](#) and select only what you want to receive.